

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave Blank)		2. REPORT DATE 29 April 99	3. REPORT TYPE AND DATES COVERED Final Report 01 Feb 1998 - 31 Dec 1998	
4. TITLE AND SUBTITLE COMPUTER FOR AB INITIO MOLECULAR DYNAMICS SIMULATIONS			5. FUNDING NUMBERS N00014-98-1-0323	
6. AUTHORS Gregory A. Voth				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry and Henry Eyring Center for Theoretical Chemistry, 315 S. 1400 E. Rm Dock, University of Utah, Salt Lake City, UT 84112			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research Ballston Centre Tower One 800 North Quincy Street, Arlington, VA 22217-5660			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) This DURIP funding was used to contribute to the expansion of an IBM SP parallel computer and its support hardware. This computer resource is being utilized for the computationally demanding ab initio molecular dynamics simulations that are part of the P.I.'s ONR funded research program. The goal of this program is to compute realistic rates for electrochemical reactions. The simulations are extraordinarily demanding in CPU time and memory requirements and have begun to yield very important new results for the water/metal electrode interface.				
14. SUBJECT TERMS			15. NUMBER OF PAGES 1	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-1
298-102

ONR Seattle

MAY 10 1999

Final Technical Report
ONR DURIP Grant N00014-98-1-0323
PR Number 98PR04001-00

Equipment Acquired by Name, Manufacturer, and Cost:

8-node SP system	IBM	\$99,000
Two Origin 200 Servers	Silicon Graphics	\$18,126
One 9mm exabyte tape drive	Ramsys	\$1,323
Two 18 Gbyte disk drives	Ramsys	\$2,176
Repair and maintenance	Silicon Graphics	<u>\$365</u>
Total		\$121,170

Special Circumstances:

The equipment purchased deviated only slightly from the original proposal in that the \$121,170 provided by the ONR DURIP grant was used to leverage and addition \$333,000 from the University of Utah to add eight, instead of four, nodes to the IBM SP system, and to double the memory on each node. One Silicon Graphics Origin200 server was purchased to help provide local data storage which could not be accommodated on the SP, and one half of another SGI Origin200 was purchased to provide graphical analysis of the data produced by the simulations on the SP. A 9mm exabyte tape drive was purchased for back-up purposes, and two 18 Gbyte disk drives were purchased for the SGI server to expand disk storage capacity.

Use of the Equipment:

The research usage of the equipment purchased conforms to that which was described in the original DURIP proposal as funded by ONR. Briefly, we are carrying out the first fully *ab initio* simulations of the water/metal interface. These simulations show that significant coupling between solvent molecules and surface states can occur even in the presence of a disordered solvent. This coupling is expected to have significant effects in heterogeneous electron transfer between the solid and aqueous redox species because the solvent-surface electronic coupling facilitates the transport of electron density away from the surface. In turn, the electron exchange between the solid and redox species is also coupled to specific solvent fluctuations in a non-trivial way. By combining the information obtained from these simulation with largescale classical molecular dynamics simulations using the appropriate effective Hamiltonians developed in our group, the accurate calculation of heterogeneous electron transfer rates for realistic systems will become a reality.

ONR Seattle

MAY 10 1999

19990518 071